

Many-body problem in Kaluza-Klein models: negative result

Alexey Chopovsky,^{1,*} Maxim Eingorn,^{2,3,†} and Alexander Zhuk^{2,‡}

¹*Department of Theoretical Physics, Odessa National University, Street Dvoryanskaya 2, Odessa 65082, Ukraine*

²*Astronomical Observatory, Odessa National University, Street Dvoryanskaya 2, Odessa 65082, Ukraine*

³*Department of Theoretical and Experimental Nuclear Physics,
Odessa National Polytechnic University, Shevchenko av. 1, Odessa 65044, Ukraine*

We consider a system of gravitating bodies in Kaluza-Klein models with toroidal compactification of extra dimensions. To simulate the ordinary astrophysical objects (e.g., our Sun), we suppose that these bodies have nonrelativistic (pressureless) equations of state in the external/our space. At the same time, they may have nonzero parameters $\omega_{(\bar{\alpha}-3)}$ ($\bar{\alpha} = 4, \dots, D$) of the equations of state in the extra dimensions. We prove that the Lagrange function of this many-body system can be constructed only if $\Sigma = \sum_{\bar{\alpha}} \omega_{(\bar{\alpha}-3)} = 0$. This strongly contradicts the observations which demand $\Sigma \approx (3 - D)/2 \sim O(1)$ for $D > 3$. Therefore, considered multidimensional Kaluza-Klein models face a severe problem.

PACS numbers: 04.25.Nx, 04.50.Cd, 04.80.Cc, 11.25.Mj

Introduction.— The idea of multidimensionality of our Universe demanded by the theories of unification of the fundamental interactions is one of the most breathtaking ideas of theoretical physics. It takes its origin from the pioneering papers by Th. Kaluza and O. Klein [1], and now the most self-consistent modern theories of unification such as superstrings, supergravity and M-theory are constructed in spacetimes with extra dimensions (see, e.g., [2]). Different aspects of the idea of multidimensionality are intensively used in numerous modern articles.

Therefore, it is important to find experimental evidence for the existence of the extra dimensions. For example, one of the aims of Large Hadronic Collider consists in detecting of Kaluza-Klein (KK) particles which correspond to excitations of the internal spaces (see, e.g., [3]).

On the other hand, if we can show that the existence of the extra dimensions is contrary to observations, then these theories are prohibited.

In our previous papers [4–6] devoted to KK models with toroidal compactification of the extra dimensions, we have shown that gravitating masses should have tension in the internal space to be in agreement with gravitational experiments in the Solar system. For example, black strings/branes with the parameter $\omega = -1/2$ of the equation of state in the internal space satisfy this condition. For this value of ω , the variations of the internal space volume are absent [7]. In the dust-like case with $\omega = 0$, such variations generate the fifth force, that leads to contradictions with the experimental data.

In the present letter we continue the investigation of the KK models with toroidal compactification. Here, we try to construct the Lagrange function for a many-body system in these models, and we come into conflict. On the one hand, the Lagrange function exists if the sum Σ of the parameters of the equations of state in the extra dimensions is equal to zero: $\Sigma = 0$. On the other hand, the agreement with the gravitational experiments

requires $\Sigma \sim O(1)$. Thus we claim that considered multidimensional KK models face a severe problem.

Nonrelativistic gravitational field for a many-body system.— To construct the Lagrange function of a system of N massive bodies in $(D + 1)$ -dimensional spacetime, we define first the nonrelativistic gravitational field created by this system. To do it, we need to get the metric coefficients in the weak field limit. The general form of the multidimensional metrics is

$$ds^2 = g_{ik} dx^i dx^k = g_{00} (dx^0)^2 + 2g_{0\mu} dx^0 dx^\mu + g_{\mu\nu} dx^\mu dx^\nu, \quad (1)$$

where the Latin indices $i, k = 0, 1, \dots, D$ and the Greek indices $\mu, \nu = 1, \dots, D$. We make the natural assumption that in the case of the absence of matter sources the spacetime is Minkowski spacetime: $g_{00} = \eta_{00} = 1$, $g_{0\mu} = \eta_{0\mu} = 0$, $g_{\mu\nu} = \eta_{\mu\nu} = -\delta_{\mu\nu}$. In our letter, we consider in detail the case where the extra dimensions have the topology of tori. In the presence of matter, the metrics is not Minkowskian one, and we investigate it in the weak field limit. It means that the gravitational field is weak and velocities of test bodies are small compared with the speed of light c . In the weak field limit the metrics is only slightly perturbed from its flat spacetime value. We will define the metrics (1) up to $1/c^2$ correction terms. Because the coordinate $x^0 = ct$, the metric coefficients can be expressed as follows:

$$g_{00} \approx 1 + h_{00} + f_{00}, \quad g_{0\mu} \approx h_{0\mu} + f_{0\mu}, \quad g_{\mu\nu} \approx -\delta_{\mu\nu} + h_{\mu\nu}, \quad (2)$$

where $h_{ik} \sim O(1/c^2)$, $f_{00} \sim O(1/c^4)$ and $f_{0\mu} \sim O(1/c^3)$. In particular, $h_{00} \equiv 2\varphi/c^2$. Later we will demonstrate that φ is the nonrelativistic gravitational potential. To get these correction terms, we should solve (in the corresponding orders of $1/c$) the multidimensional Einstein

equation

$$R_{ik} = \frac{2S_D \tilde{G}_D}{c^4} \left(T_{ik} - \frac{1}{D-1} g_{ik} T \right), \quad (3)$$

where $S_D = 2\pi^{D/2}/\Gamma(D/2)$ is the total solid angle (the surface area of the $(D-1)$ -dimensional sphere of the unit radius), $\tilde{G}_D$ is the gravitational constant in the $(D = D+1)$ -dimensional spacetime. We consider a system of N discrete massive (with rest masses m_p , $p = 1, \dots, N$) bodies. We suppose that the pressure of these bodies in the external three-dimensional space is much less than their energy density. This is a natural approximation for ordinary astrophysical objects such as our Sun. For example, this approach works well for calculating the gravitational experiments in the Solar system [8]. In other words, the gravitating bodies are pressureless in the external/our space. On the other hand, we suppose that they may have pressure in the extra dimensions. Therefore, nonzero components of the energy-momentum tensor of the system can be written in the following form:

$$T^{ik} = \tilde{\rho} c^2 u^i u^k, \quad i, k = 0, \dots, 3, \quad (4)$$

$$T^{i\bar{\alpha}} = \tilde{\rho} c^2 u^i u^{\bar{\alpha}}, \quad i = 0, \dots, 3; \bar{\alpha} = 4, \dots, D, \quad (5)$$

$$T^{\bar{\alpha}\bar{\beta}} = -p_{(\bar{\alpha}-3)} g^{\bar{\alpha}\bar{\beta}} + \tilde{\rho} c^2 u^{\bar{\alpha}} u^{\bar{\beta}}, \quad \bar{\alpha}, \bar{\beta} = 4, \dots, D \quad (6)$$

where the $(D+1)$ -velocity $u^i = dx^i/ds$ and

$$\tilde{\rho} \equiv \sum_{p=1}^N [(-1)^D g]^{-1/2} m_p \sqrt{g_{lm} \frac{dx^l}{dx^0} \frac{dx^m}{dx^0}} \delta(\mathbf{x} - \mathbf{x}_p), \quad (7)$$

where $\mathbf{x}_p$ is a D -dimensional radius-vector of the p -th particle. In what follows, the Greek indices $\alpha, \beta = 1, 2, 3$; $\bar{\alpha}, \bar{\beta} = 4, \dots, D$ and μ, ν still run from 1 to D . In the extra dimensions we suppose the equations of state:

$$p_{(\bar{\alpha}-3)} = \omega_{(\bar{\alpha}-3)} \tilde{\rho} c^2. \quad (8)$$

If all parameters $\omega_{(\bar{\alpha}-3)} = 0$, then we come back to the model considered in our paper [4]. Here, massive bodies have dust-like equations of state in all spatial dimensions. If all $\omega_{(\bar{\alpha}-3)} = -1/2$ (tension in the extra dimensions), then these equations of state correspond to black strings (in the case of one extra dimension, i.e. $D = 4$) and black branes (for $D > 4$). If parameters satisfy the condition $\sum_{\bar{\alpha}} \omega_{(\bar{\alpha}-3)} \equiv \Sigma = -(D-3)/2$, then this case corresponds to latent solitons [6]. Obviously, black strings/branes satisfy this condition.

Now, we will solve the Einstein equation (3) in the same way as it was done in [4]. Obviously, for $\omega_{(\bar{\alpha}-3)} = 0$, $\bar{\alpha} = 4, \dots, D$, we should reproduce the results of this paper. Up to $O(1/c^2)$, we get the following nonzero components

$$h_{00} = \frac{2\varphi}{c^2}, \quad h_{\alpha\beta} = \frac{1-\Sigma}{D-2+\Sigma} \frac{2\varphi}{c^2} \delta_{\alpha\beta}, \quad (9)$$

$$h_{\bar{\alpha}\bar{\beta}} = \frac{\omega_{(\bar{\alpha}-3)}(D-1) + 1 - \Sigma}{D-2+\Sigma} \frac{2\varphi}{c^2} \delta_{\bar{\alpha}\bar{\beta}}, \quad (10)$$

where φ satisfies the D -dimension Poisson equation

$$\Delta_D \varphi(\mathbf{x}) = S_D \tilde{G}_D \frac{2(D-2+\Sigma)}{D-1} \rho(\mathbf{x}). \quad (11)$$

Here, $\Delta_D = \delta^{\mu\nu} \partial^2 / \partial x^\mu \partial x^\nu$ and the rest mass density is

$$\rho(\mathbf{x}) = \sum_{p=1}^N m_p \delta(\mathbf{x} - \mathbf{x}_p). \quad (12)$$

To get the solutions (9) and (10), we use the standard (see, e.g., Eq. (105.10) in [8]) gauge condition

$$\partial_k \left(h_i^k - \frac{1}{2} h_l^l \delta_i^k \right) = 0. \quad (13)$$

It can be easily verified that this condition is satisfied (up to $O(1/c^2)$) for $i = 0, \alpha$. However, for $i = \bar{\alpha}$, we should demand either $\omega_{(\bar{\alpha}-3)} = 0$ or $\partial_{\bar{\alpha}} \varphi = 0$. Because we consider the general case $\omega_{(\bar{\alpha}-3)} \neq 0$, we must choose the latter condition. Therefore, the presence of nonzero pressure/tension in the extra dimensions results in the metric coefficients which do not depend on the coordinates of the internal space, i.e. the gravitating masses should be uniformly smeared over the extra dimensions. In this case, the rest mass density (12) should be rewritten in the form: $\rho(\mathbf{x}) \rightarrow \rho(\mathbf{r}) = \sum_p m_p \delta(\mathbf{r} - \mathbf{r}_p) / \prod_{\bar{\alpha}} a_{(\bar{\alpha}-3)}$, where $\mathbf{r}_p$ is a three-dimensional radius vector of the p -th particle in the external space, $a_{(\bar{\alpha}-3)}$ are periods of the tori (i.e. $\prod_{\bar{\alpha}} a_{(\bar{\alpha}-3)}$ is the volume of the internal space). Then, Eq. (11) is reduced to the ordinary three-dimensional Poisson equation

$$\Delta_3 \varphi(\mathbf{r}) = 4\pi G_N \sum_p m_p \delta(\mathbf{r} - \mathbf{r}_p) \quad (13)$$

with the solution

$$\varphi(\mathbf{r}) = - \sum_p \frac{G_N m_p}{|\mathbf{r} - \mathbf{r}_p|}, \quad (14)$$

where G_N is the Newtonian gravitational constant:

$$4\pi G_N = \frac{2S_D(D-2+\Sigma)}{(D-1) \prod_{\bar{\alpha}} a_{(\bar{\alpha}-3)}} \tilde{G}_D. \quad (15)$$

Following the paper [4], we can also obtain the $O(1/c^3)$ and $O(1/c^4)$ correction terms. As a result, the metric coefficients read

$$\begin{aligned} g_{00} \approx & 1 + \frac{2\varphi(\mathbf{r})}{c^2} + \frac{2\varphi^2(\mathbf{r})}{c^4} + \frac{(D-2)^2 - \Sigma D}{(D-2+\Sigma)(D-2)} \frac{2G_N^2}{c^4} \\ & \times \sum_p \frac{m_p}{|\mathbf{r} - \mathbf{r}_p|} \sum_{q \neq p} \frac{m_q}{|\mathbf{r}_p - \mathbf{r}_q|} \\ & - \frac{D-\Sigma}{D-2+\Sigma} \frac{G_N}{c^4} \sum_p \frac{m_p v_p^2}{|\mathbf{r} - \mathbf{r}_p|}, \end{aligned} \quad (16)$$

$$g_{0\alpha} \approx \frac{3D-2-\Sigma}{D-2+\Sigma} \frac{G_N}{2c^3} \sum_p \frac{m_p}{|\mathbf{r}-\mathbf{r}_p|} v_p^\alpha + \frac{G_N}{2c^3} \sum_p \frac{m_p}{|\mathbf{r}-\mathbf{r}_p|} n_p^\alpha (\mathbf{n}_p \mathbf{v}_p), \quad (17)$$

$$g_{\alpha\beta} \approx \left(-1 + \frac{1-\Sigma}{D-2+\Sigma} \frac{2\varphi(\mathbf{r})}{c^2} \right) \delta_{\alpha\beta}, \quad (18)$$

$$g_{\bar{\alpha}\bar{\beta}} \approx \left(-1 + \frac{\omega_{(\bar{\alpha}-3)}(D-1) + 1 - \Sigma 2\varphi(\mathbf{r})}{D-2+\Sigma} \frac{1}{c^2} \right) \delta_{\bar{\alpha}\bar{\beta}}. \quad (19)$$

Here, the three-velocity $v_p^\alpha = dx_p^\alpha/dt$, the three-dimensional unit vector $n_p^\alpha = (x^\alpha - x_p^\alpha)/|\mathbf{r} - \mathbf{r}_p|$ and $(\mathbf{n}_p \mathbf{v}_p) = \sum_\alpha n_p^\alpha v_p^\alpha$. Obviously, if $\omega_{(\bar{\alpha}-3)} = 0$, $\forall \bar{\alpha}$, $\Rightarrow \Sigma = 0$, these formulas are reduced to ones in [4], which in turn coincide with the metric coefficients obtained in [8] in the case $D = 3$.

Lagrange function for a two-body system.— Let us construct now the Lagrange function of the many-body system described above. To perform it we will follow the procedure described in [8] (see §106). The Lagrange function of a particle a with the mass m_a in the gravitational field created by the other bodies is given by the expression

$$L_a = -m_a c \frac{ds_a}{dt} \quad (20)$$

$$= -m_a c^2 \left(g_{00} + 2 \sum_\mu g_{0\mu} \frac{v_a^\mu}{c} + \sum_{\mu\nu} g_{\mu\nu} \frac{v_a^\mu v_a^\nu}{c^2} \right)^{1/2},$$

where the metric coefficients are taken at $\mathbf{r} = \mathbf{r}_a$. For our purposes, it is sufficient to consider the case of two particles. The substitution of the metric coefficients (16)-(19) leads to the following expression for the particle "1":

$$L_1 = f(\mathbf{v}_1^2) + G_N \frac{m_1 m_2}{|\mathbf{r} - \mathbf{r}_2|} - \frac{1}{2c^2} G_N^2 \frac{m_1 m_2^2}{|\mathbf{r} - \mathbf{r}_2|^2} - \frac{1}{c^2} \mathbf{a}(D, \Sigma) G_N^2 \frac{m_1^2 m_2}{|\mathbf{r} - \mathbf{r}_2| |\mathbf{r}_1 - \mathbf{r}_2|} + \frac{1}{2c^2} \frac{G_N m_1 m_2}{|\mathbf{r} - \mathbf{r}_2|} \left[\mathbf{b}(D, \Sigma) v_2^2 + 2 \left(\mathfrak{d}(D, \Sigma) + \frac{1}{2} \right) v_1^2 - \mathbf{c}(D, \Sigma) (\mathbf{v}_1 \mathbf{v}_2) - (\mathbf{n}_2 \mathbf{v}_2) (\mathbf{n}_2 \mathbf{v}_1) \right], \quad (21)$$

where $f(\mathbf{v}_1^2) = m_1 v_1^2/2 + m_1 v_1^4/(8c^2)$ and we drop the term $-m_1 c^2$. Here, we use the following abbreviations:

$$\mathbf{a}(D, \Sigma) \equiv \frac{(D-2)^2 - \Sigma D}{(D-2+\Sigma)(D-2)}, \quad \mathbf{b}(D, \Sigma) \equiv \frac{D-\Sigma}{D-2+\Sigma},$$

$$\mathbf{c}(D, \Sigma) \equiv \frac{3D-2-\Sigma}{D-2+\Sigma}, \quad \mathfrak{d}(D, \Sigma) = \frac{1-\Sigma}{D-2+\Sigma}. \quad (22)$$

The total Lagrange function of the two-body system should be constructed so that it leads to the correct values of the forces $\partial L_a / \partial \mathbf{r}|_{\mathbf{r}=\mathbf{r}_a}$ acting on each of the bodies for given motion of the others [8]. Following this prescription, we obtain from (21) the two-body Lagrange function

$$L_1^{(2)} = \tilde{f}(\mathbf{v}_1^2, \mathbf{v}_2^2) + \frac{G_N m_1 m_2}{r_{12}} - \frac{G_N^2 m_1 m_2 (\mathbf{a} m_1 + m_2)}{2c^2 r_{12}^2} + \frac{G_N m_1 m_2}{2c^2 r_{12}} [\mathbf{b}(D, \Sigma) v_2^2 + (2\mathfrak{d}(D, \Sigma) + 1) v_1^2 - \mathbf{c}(D, \Sigma) (\mathbf{v}_1 \mathbf{v}_2) - (\mathbf{n}_{12} \mathbf{v}_1) (\mathbf{n}_{12} \mathbf{v}_2)], \quad (23)$$

where $\tilde{f}(\mathbf{v}_1^2, \mathbf{v}_2^2) = \sum_{a=1}^2 m_a v_a^2/2 + \sum_{a=1}^2 m_a v_a^4/(8c^2)$. It can be easily seen that $\partial L_1 / \partial \mathbf{r}|_{\mathbf{r}=\mathbf{r}_1} = \partial L_1^{(2)} / \partial \mathbf{r}_1$. By the same way we can construct the two-body Lagrange function $L_2^{(2)}$ from the Lagrange function L_2 for the particle "2":

$$L_2^{(2)} = \tilde{f}(\mathbf{v}_1^2, \mathbf{v}_2^2) + \frac{G_N m_1 m_2}{r_{12}} - \frac{G_N^2 m_1 m_2 (m_1 + \mathbf{a} m_2)}{2c^2 r_{12}^2} + \frac{G_N m_1 m_2}{2c^2 r_{12}} [\mathbf{b}(D, \Sigma) v_1^2 + (2\mathfrak{d}(D, \Sigma) + 1) v_2^2 - \mathbf{c}(D, \Sigma) (\mathbf{v}_1 \mathbf{v}_2) - (\mathbf{n}_{12} \mathbf{v}_1) (\mathbf{n}_{12} \mathbf{v}_2)]. \quad (24)$$

It is worth noting that both $L_1^{(2)}$ and $L_2^{(2)}$ are reduced to the Lagrange function of the two-body system in [8] if we assume that $D = 3$, $\Sigma = 0$.

Obviously, the Lagrange functions $L_1^{(2)}$ and $L_2^{(2)}$ should be symmetric with respect to permutations of particles 1 and 2 and should coincide with each other. This requires the following conditions:

$$\mathbf{a}(D, \Sigma) = 1, \quad \mathbf{b}(D, \Sigma) = 2\mathfrak{d}(D, \Sigma) + 1. \quad (25)$$

The second condition is satisfied identically. However, the former one is valid only for

$$\Sigma = 0. \quad (26)$$

This happens, e.g., for the dust-like equations of state in the extra dimensions: $\omega_{(\bar{\alpha}-3)} = 0$, $\forall \bar{\alpha}$. Definitely, black strings/branes contradict this condition.

We can also get an additional restriction on Σ from the known gravitational experiments in the Solar system. For example, the post-Newtonian parameter γ for the considered models is expressed via Σ as follows (see, e.g., [4–6]):

$$\gamma - 1 = \frac{3 - D - 2\Sigma}{D - 2 + \Sigma}. \quad (27)$$

There are strong experimental restrictions on the value of γ . The tightest constraint on γ comes from the Shapiro time-delay experiment using the Cassini spacecraft, namely: $\gamma - 1 = (2.1 \pm 2.3) \times 10^{-5}$ [9–11]. Therefore, $\gamma - 1 \approx 0 \Rightarrow \Sigma \approx (3 - D)/2 \sim O(1)$ for $D > 3$. It is worth

noting that for latent solitons (and black strings/branes as their particular cases) $\Sigma = (3 - D)/2$ [6] and $\gamma = 1$, as in general relativity. This is in good agreement with observations.

Hence, we come into conflict. On the one hand, the Lagrange function for a two-body system exists for $\Sigma = 0$. On the other hand, this value contradicts the gravitational experiments.

Summary.— In this letter, we have constructed the Lagrange function for a two-body system in the case of Kaluza-Klein models with toroidal compactification of the extra dimensions. The case of more than two bodies is straightforward. We supposed that gravitating bodies are pressureless in the external/our space. This is a natural approximation for ordinary astrophysical objects such as our Sun. For example, this approach works well for calculating the gravitational experiments in the Solar system [8]. On the other hand, we suppose that gravitating masses may have nonzero parameters $\omega_{(\bar{\alpha}-3)}$ ($\bar{\alpha} = 4, \dots, D$) of the equations of state in the extra dimensions. We prove that the Lagrange function of this many-body system can be constructed only if $\Sigma = \sum_{\bar{\alpha}} \omega_{(\bar{\alpha}-3)} = 0$. This strongly contradicts the observations which demand $\Sigma \approx (3 - D)/2 \sim O(1)$ for $D > 3$. Hence, we come into conflict.

It is not difficult to realize that this result can be generalized to the case of arbitrary Ricci-flat internal spaces. Calabi-Yau manifolds are of special interest because these Ricci-flat spaces are widely used in superstring theories. We will elaborate this general case in our forthcoming paper.

One more important result obtained in this latter is worth noting. It is well known [5–7] that tension in the internal spaces is the necessary condition to satisfy the gravitational experiments in KK models with toroidal compactification. In our letter, we have shown that the presence of pressure/tension in the internal space leads necessarily to the uniform smearing of the gravitating masses over the internal space. For example, black strings/branes have tension in the internal space (see, e.g., [12]). Therefore, they should be smeared. However, uniformly smeared gravitating bodies cannot have

excited KK states (KK particles), which looks unnatural from the point of quantum mechanics.

Therefore, considered multidimensional Kaluza-Klein models face severe problems.

This work was supported in part by the "Cosmomicrophysics-2" programme of the Physics and Astronomy Division of the National Academy of Sciences of Ukraine.

* Electronic address: alexey.chopovsky@gmail.com

† Electronic address: maxim.eingorn@gmail.com

‡ Electronic address: ai.zhuk2@gmail.com

- [1] Th. Kaluza, *Zum Unitätsproblem der Physik*, Sitzungsber. d. Preuss. Akad. d. Wiss., 966 (1921); O. Klein, *Quantentheorie und fünfdimensionale Relativitätstheorie*, Zeitschrift für Physik **37**, 895 (1926).
- [2] J. Polchinski, *String Theory, Volume 2: Superstring Theory and Beyond* (Cambridge University Press, Cambridge, 1998).
- [3] G. Bhattacharyya, A. Datta, S.K. Majee and A. Raychaudhuri, Nucl. Phys. B **821**, 48 (2009); arXiv:hep-ph/0904.0937.
- [4] M. Eingorn and A. Zhuk, Class. Quant. Grav. **27**, 205014 (2010); arXiv:gr-qc/1003.5690.
- [5] M. Eingorn and A. Zhuk, Phys. Rev. D **83**, 044005 (2011); arXiv:qr-qc/1010.5740.
- [6] M. Eingorn, O. de Medeiros, L. Crispino and A. Zhuk, Phys. Rev. D **84**, 024031 (2011); arXiv:gr-qc/1101.3910.
- [7] M. Eingorn and A. Zhuk, Phys. Lett. B **713**, 154-159 (2012); arXiv:gr-qc/1201.1756.
- [8] L.D. Landau and E.M. Lifshitz, *The Classical Theory of Fields, Fourth Edition: Volume 2 (Course of Theoretical Physics Series)* (Pergamon Press, Oxford, 2000).
- [9] C.M. Will, *Was Einstein Right? Testing Relativity at the Centenary*. In *100 Years of Relativity: Space-time Structure: Einstein and Beyond*, edited by A. Ashtekar (World Scientific, Singapore, 2005); arXiv:gr-qc/0504086.
- [10] Bh. Jain and J. Khoury, *Cosmological Tests of Gravity* (2010); arXiv:astro-ph/1004.3294.
- [11] B. Bertotti, L. Iess, and P. Tortora, Nature **425**, 374 (2003).
- [12] D. Kastor and J. Traschen, JHEP **0609**, 022 (2006); arXiv:hep-th/0607051.